

University of Groningen

Efficacy of tiotropium in adults with moderate asthma, by leukotriene receptor antagonist use at baseline

Hashimoto, Shu; Casale, Thomas B; Engel, Michael; Moroni-Zentgraf, Petra; Bour, Louis J; Kerstjens, Huib A M

Published in:

Allergology international : official journal of the Japanese Society of Allergology

DOI:

[10.1016/j.alit.2017.12.002](https://doi.org/10.1016/j.alit.2017.12.002)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hashimoto, S., Casale, T. B., Engel, M., Moroni-Zentgraf, P., Bour, L. J., & Kerstjens, H. A. M. (2018). Efficacy of tiotropium in adults with moderate asthma, by leukotriene receptor antagonist use at baseline. *Allergology international : official journal of the Japanese Society of Allergology*, 67(3), 411-413. <https://doi.org/10.1016/j.alit.2017.12.002>

Copyright


Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



Letter to the Editor

Efficacy of tiotropium in adults with moderate asthma, by leukotriene receptor antagonist use at baseline

Dear Editor,

In Japan and the United States, leukotriene receptor antagonists (LTRAs) are commonly prescribed for controlling asthma.^{1,2} The Japanese guidelines for adult asthma recommend LTRAs for adult patients with persistent asthma as an add-on controller option to ICSs or a combination of ICS and LABAs.³ Moreover, the co-administration of ICS and LTRAs creates an additional bronchodilator effect in asthma patients.⁴

Tiotropium, a long-acting muscarinic antagonist (LAMA), has been newly included in the 2016 Global Initiative for Asthma (GINA) guidelines as a bronchodilator treatment for asthma.⁵ In phase 3 trials replicating the MezzoTinA trial, tiotropium had similar efficacy and tolerability to salmeterol when given as add-on treatment to medium-dose ICS, with or without LTRAs, in adults with persistent asthma.⁶ Around 9% of patients in the MezzoTinA

trials were symptomatic at baseline despite treatment with both ICS and LTRAs. Here, we examine whether LTRA use at baseline impacts the efficacy of tiotropium as add-on treatment to medium-dose ICS in a post hoc analysis of data from the MezzoTinA trials.

Detailed methods have been reported previously.⁶ Briefly, eligible patients were between 18 and 75 years of age, with symptomatic asthma (mean Asthma Control Questionnaire [ACQ-7] score of ≥ 1.5), a pre-bronchodilator FEV₁ 60–90% of predicted normal, were under treatment with medium-dose ICS (400–800 µg budesonide or equivalent) and had never smoked or had a smoking history of <10 pack-years, with no smoking 1 year before enrolment. Patients with COPD or who used concomitant LAMAs or LABAs within 4 weeks prior to randomisation were excluded. Patients could continue use of LTRAs if taking them within 3 months before screening.

Table 1
Demographics and baseline patient characteristics.

	LTRA use at baseline							
	No (n = 1917)				Yes (n = 183)			
	Tiotropium 5 µg QD (n = 472)	Tiotropium 2.5 µg QD (n = 471)	Salmeterol 50 µg BID (n = 494)	Placebo [†] (n = 480)	Tiotropium 5 µg QD (n = 45)	Tiotropium 2.5 µg QD (n = 48)	Salmeterol 50 µg BID (n = 47)	Placebo [†] (n = 43)
Age, years	44.4 ± 12.8	43.2 ± 12.7	42.3 ± 12.7	42.4 ± 12.9	43.6 ± 11.0	44.9 ± 14.1	40.3 ± 14.4	47.0 ± 13.4
Female, n (%)	275 (58.3)	280 (59.4)	279 (56.5)	283 (59.0)	25 (55.6)	36 (75.0)	33 (70.2)	28 (65.1)
Race, n (%)								
American Indian/Alaska Native	27 (5.7)	26 (5.5)	30 (6.1)	28 (5.8)	3 (6.7)	1 (2.1)	1 (2.1)	2 (4.7)
Asian	205 (43.4)	195 (41.4)	216 (43.7)	199 (41.5)	20 (44.4)	25 (52.1)	13 (27.7)	20 (46.5)
Black/African American	19 (4.0)	16 (3.4)	13 (2.6)	26 (5.4)	3 (6.7)	1 (2.1)	2 (4.3)	1 (2.3)
Hawaiian/Pacific Islander	0 (0.0)	2 (0.4)	0 (0.0)	1 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
White	221 (46.8)	232 (49.3)	235 (47.6)	226 (47.1)	19 (42.2)	21 (43.8)	31 (66.0)	20 (46.5)
Body mass index, kg/m ²	26.9 ± 6.1	26.6 ± 6.2	26.6 ± 6.2	27.0 ± 6.3	29.0 ± 7.8	26.7 ± 5.6	27.6 ± 6.8	26.7 ± 5.8
Smoking status, n (%)								
Ex-smoker	87 (18.4)	71 (15.1)	77 (15.6)	66 (13.8)	10 (22.2)	11 (22.9)	18 (38.3)	4 (9.3)
Never smoked	385 (81.6)	400 (84.9)	417 (84.4)	414 (86.3)	35 (77.8)	37 (77.1)	29 (61.7)	39 (90.7)
Smoking history, pack-years	4.3 ± 2.8	4.0 ± 2.9	4.2 ± 2.7	4.0 ± 2.4	6.3 ± 4.6	3.8 ± 2.8	3.5 ± 2.8	6.1 ± 3.9
Duration of asthma, years	22.9 ± 15.1	22.0 ± 14.6	20.7 ± 14.0	20.8 ± 13.5	24.0 ± 14.2	22.3 ± 10.9	23.2 ± 17.0	24.5 ± 14.8
ICS maintenance dose, µg [‡]	663.1 ± 219.4	654.3 ± 216.8	646.4 ± 207.9	665.9 ± 216.5	672.0 ± 177.6	671.7 ± 175.2	696.2 ± 170.0	695.8 ± 227.3
FEV ₁ , mL, at randomisation (pre bronchodilation)	2203 ± 630	2269 ± 649	2327 ± 655	2275 ± 666	2224 ± 635	2227 ± 670	2429 ± 678	2083 ± 706
FEV ₁ , % predicted normal	74.1 ± 11.2	75.2 ± 11.5	75.7 ± 11.7	75.2 ± 11.4	72.3 ± 12.3	77.4 ± 11.7	77.5 ± 11.0	73.1 ± 12.6
ACQ-7 score	2.2 ± 0.5	2.2 ± 0.5	2.2 ± 0.5	2.2 ± 0.5	2.2 ± 0.5	2.2 ± 0.5	2.2 ± 0.4	2.2 ± 0.5
AQLQ score	4.8 ± 1.0	4.8 ± 0.9	4.9 ± 0.9	4.9 ± 0.9	4.8 ± 1.0	4.8 ± 0.9	4.8 ± 0.9	4.7 ± 0.8

Treated set. All values are mean ± SD unless otherwise stated.

ACQ-7, Asthma Control Questionnaire; AQLQ, Asthma Quality of Life Questionnaire; BID, twice daily; LTRA, leukotriene receptor antagonist; QD, once daily.

[†] Placebo Respimat[®] once daily plus placebo hydrofluoroalkane metered-dose inhaler twice daily.

[‡] Budesonide 400–800 µg or equivalent dose.

Peer review under responsibility of Japanese Society of Allergy.

<https://doi.org/10.1016/j.alit.2017.12.002>

1323-8930/Copyright © 2018, Japanese Society of Allergy. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

After screening and a 4-week run-in period, patients were randomised (1:1:1:1) to self-administer once-daily tiotropium 5 µg via Respimat® inhaler; once-daily tiotropium 2.5 µg (same device); twice-daily salmeterol 50 µg via hydrofluoroalkane metered-dose inhaler (HFA-MDI); or placebo (double-dummy design). The study drug was administered as an add-on to maintenance treatment with ICS (400–800 µg budesonide or equivalent). Rescue treatment (salbutamol HFA-MDI) was permitted throughout the study. For acute asthma exacerbations, temporary ICS dose increases and temporary addition of systemic corticosteroids or theophylline preparations were allowed.

Spirometric lung function tests were conducted to determine peak and trough FEV₁ values at randomisation and throughout the treatment period. Peak FEV₁ response, measured within 3 h after dosing (FEV_{1(0–3 h)}), and trough FEV₁ response, measured at the end of the dosing interval (24 h after previous dose and 10 min before next dose), at the end of the 24-week treatment period were stratified by baseline LTRA use (yes/no).

Patients participated from 233 sites in 14 countries (Latvia, Poland, Romania, Russia, Brazil, China, Colombia, Germany, Guatemala, India, Japan, Mexico, Peru, and the United States). Of 2100 randomised patients (Japanese, 240), 183 were using an LTRA at baseline (of these, tiotropium 5 µg was administered to 45 patients; tiotropium 2.5 µg, 48; salmeterol 50 µg, 47; placebo, 43). Overall, 1975 patients completed the study. Demographics and baseline patient characteristics were comparable regardless of LTRA use at baseline (Table 1). While only a slightly higher proportion of Japanese patients with asthma had comorbid allergic rhinitis compared to the overall population (81/240 [33.8%] vs. 608/2100 [29.0%], respectively), the proportion of Japanese patients taking LTRAs was 3-fold higher than the overall population (60/240 [25.0%] vs. 183/2100 [8.7%], respectively).

Peak FEV_{1(0–3 h)} and trough FEV₁ improvements were independent of LTRA use at baseline (interaction p values = 0.9057 and 0.9083, respectively) (Fig. 1). Compared with the placebo group, adjusted mean differences for peak FEV_{1(0–3 h)} increased




Fig. 1. Differences in lung function at week 24 by LTRA use at baseline compared with placebo; (A) Peak FEV_{1(0–3 h)}, (B) Trough FEV₁. Full analysis set adjusted for treatment, study, visit, baseline, treatment-by-visit and baseline-by-visit. The interaction term 'treatment-by-subgroup' and the term 'subgroup' were added to the linear model to evaluate the treatment effect across the subgroups. Common baseline mean ± SD = 2265 ± 653 mL. Subgroup baseline values: No = 2267 mL; Yes = 2244 mL. Number of patients receiving placebo[†]: overall, n = 492; LTRA use at baseline: No, n = 454; LTRA use at baseline: Yes, n = 38. [†]Placebo Respimat® once daily plus placebo HFA-MDI twice daily. FEV_{1(0–3 h)}, FEV₁ measured within 3 h of dosing; LTRA, leukotriene receptor antagonist.

significantly in both patients taking an LTRA at baseline ($p < 0.05$) and in patients not taking an LTRA at baseline ($p < 0.0001$). Adjusted mean differences for trough FEV₁ also increased; a significant ($p < 0.0001$) difference was observed among patients not taking an LTRA at baseline, and a similar, though not significant, difference was seen in patients taking an LTRA, probably due to lower patient numbers. It must be noted that <10% of the study population used LTRAs at baseline.

As the proportion of Japanese patients taking an LTRA was 3-fold higher than the overall population, we reported peak FEV_{1(0–3 h)} and trough FEV₁ in both populations. We did not observe any difference in the improvement of respiratory function regardless of whether patients were Japanese or non-Japanese (interaction p values; peak FEV₁, $p = 0.7352$ and trough FEV₁, $p = 0.5234$; [Supplementary Table 1](#)). In addition, the proportion of ACQ-7 responders was recorded in patients with and without LTRA use at baseline. There were more ACQ-7 responders in the tiotropium 5 µg (with LTRA, OR [95% CI], 2.48 [1.00, 6.18], $p = 0.0834$; without LTRA, 1.25 [0.96, 1.63], $p = 0.1062$) and 2.5 µg (with LTRA, 1.38 [0.59, 3.22], $p = 0.5226$; without LTRA, 1.32 [1.01, 1.72], $p = 0.0456$) groups and the salmeterol group (with LTRA, 1.68 [0.71, 3.97], $p = 0.3262$; without LTRA, 1.44 [1.11, 1.87], $p = 0.0080$) than in the placebo group, regardless of LTRA use at baseline. It must be noted that the OR of ACQ-7 responders between the active drug groups and the placebo group exceeded 1 regardless of LTRA use at baseline.

Different mechanisms of action between LTRAs and tiotropium may explain these observations. While LTRAs block leukotriene D₄, which induces bronchoconstriction and inflammation associated with asthma,⁷ tiotropium binds to muscarinic M3 receptors, blocking the effects of acetylcholine on airway smooth muscle and providing 24-h bronchodilation as it slowly dissociates.⁸ Co-administration of LTRAs and ICS confers complementary effects on inflammatory markers⁹ and yields an augmented bronchodilator effect.⁴ In addition to blocking M3 muscarinic receptors on airway smooth muscle, tiotropium in combination with corticosteroids has synergistically modulated airway inflammation in an animal model of chronic asthma.¹⁰ These findings suggest that both LTRAs and tiotropium possess bronchodilator effects that are distinct and might be additive.

In conclusion, once-daily tiotropium added to ICS improves lung function in patients with moderate symptomatic asthma, independent of LTRA use at baseline. Future clinical studies with larger populations evaluating impact of LTRAs on other markers should be considered.

Acknowledgements

This study was funded by Boehringer Ingelheim. The sponsor of the study had a role in study design, data collection, data analysis, data interpretation, and writing of the report. Medical writing services and editorial support, supported financially by Boehringer Ingelheim, were provided by Ruhi Ubale, PhD, of Cactus Communications. Editorial support was provided by Yasuhiro Kurotori and Daisuke Kuroki of Nippon Boehringer Ingelheim. The authors thank Emilio Pizzichini for reviewing this manuscript.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.alit.2017.12.002>.

Conflict of interest

SH received honoraria from Boehringer Ingelheim. TBC received research grants from Boehringer Ingelheim. HAMK received research grants from Boehringer Ingelheim and Novartis; fees for patient recruitment in clinical trials from several pharmaceutical companies; and served on the advisory boards for Boehringer Ingelheim, AstraZeneca, Chiesi, GlaxoSmithKline, and Novartis. ME, PMZ, and LJB are employees of Boehringer Ingelheim.

Shu Hashimoto^{a,*}, Thomas B. Casale^b, Michael Engel^c,
Petra Moroni-Zentgraf^c, Louis J. Bour^d, Huib A.M. Kerstjens^e

^a Department of Internal Medicine, Nihon University School of Medicine, Tokyo, Japan

^b Division of Allergy and Immunology, University of South Florida Morsani College of Medicine, Tampa, FL, USA

^c TA Respiratory Diseases, Boehringer Ingelheim Pharma GmbH & Co. KG, Ingelheim am Rhein, Germany

^d Biostatistics, Boehringer Ingelheim Pharma GmbH & Co. KG, Biberach an der Riss, Germany

^e University of Groningen, Department of Pulmonary Medicine, University Medical Center Groningen, Groningen, The Netherlands

* Corresponding author. Division of Respiratory Medicine, Department of Internal Medicine, Nihon University School of Medicine, Tokyo, Japan.
E-mail address: hashimoto.shu@nihon-u.ac.jp (S. Hashimoto).

References

- Butler MG, Zhou EH, Zhang F, Wu YT, Wu AC, Levenson MS, et al. Changing patterns of asthma medication use related to US Food and Drug Administration long-acting β_2 -agonist regulation from 2005–2011. *J Allergy Clin Immunol* 2016;**137**:710–7.
- Ichinose M, Sugiura H, Nagase H, Yamaguchi M, Inoue H, Sagara H, et al. Japanese guidelines for adult asthma 2017. *Allergol Int* 2017;**66**:163–89.
- Ohta K, Ichinose M, Nagase H, Yamaguchi M, Sugiura H, Tohda Y, et al. Japanese guideline for adult asthma 2014. *Allergol Int* 2014;**63**:293–333.
- Lavolette M, Malmstrom K, Lu S, Chervinsky P, Pujet JC, Peszek I, et al. Montelukast added to inhaled beclomethasone in treatment of asthma. Montelukast/Beclomethasone Additivity Group. *Am J Respir Crit Care Med* 1999;**160**:1862–8.
- Global Initiative for Asthma (GINA). *Pocket Guide for Asthma Management and Prevention: For Adults and Children Older than 5 Years*. CreateSpace Independent Publishing Platform; 2016.
- Kerstjens HA, Casale TB, Bleecker ER, Meltzer EO, Pizzichini E, Schmidt O, et al. Tiotropium or salmeterol as add-on therapy to inhaled corticosteroids for patients with moderate symptomatic asthma: two replicate, double-blind, placebo-controlled, parallel-group, active-comparator, randomised trials. *Lancet Respir Med* 2015;**3**:367–76.
- Spector SL. Leukotrienes and the mechanism by which the LTRAs block their asthma-related effects. *Postgrad Med* 2000;**108**:6–11.
- Gosens R, Zaagsma J, Meurs H, Halayko AJ. Muscarinic receptor signaling in the pathophysiology of asthma and COPD. *Respir Res* 2006;**7**:73.
- Currie GP, Lee DK, Haggart K, Bates CE, Lipworth BJ. Effects of montelukast on surrogate inflammatory markers in corticosteroid-treated patients with asthma. *Am J Respir Crit Care Med* 2003;**167**:1232–8.
- Kistemaker LE, Bos IS, Menzen MH, Maarsingh H, Meurs H, Gosens R. Combination therapy of tiotropium and ciclesonide attenuates airway inflammation and remodeling in a Guinea pig model of chronic asthma. *Respir Res* 2016;**17**:13.

Received 14 June 2017

Received in revised form 15 November 2017

Accepted 17 November 2017

Available online 15 February 2018